

Branes in $N = 2$, $D = 4$ supergravity and the conformal field theory limit¹

Klaus Behrndt

Institut für Physik, Humboldt-Universität,
Invalidenstr. 110, D-10115 Berlin

Abstract: In this article we summarise the brane solutions (instantons, black holes and strings) in 4 dimensions when embedded in $N = 2$ supergravity. Like in 10 dimensions these solutions are related by duality transformations (T -duality, c -map). For the case that the graviphoton has no magnetic charge, we discuss the conformal field theory description of the near-horizon geometry. The decoupling of the massless modes in the matrix limit of the M -theory compactified on a Calabi-Yau threefold is discussed.

In the last years we made substantial progress in the understanding of $N=2$ black holes in 4 and 5 dimensions [1, 2, 3, 4, 5]. In order to obtain these black holes one has to compactify the 10-dimensional supergravity on an internal space with unknown metric, $K3$ for heterotic and Calabi-Yau threefold (CY_3) on type II side. As consequence, one cannot obtain them simply by dimensional reduction and hence one has to solve either the 4-d equations of motion or the Killing spinor equations.

Considering only the ungauged bosonic part the equations of motion follow from the $N=2$ supergravity action given by

$$S = \int d^4x \sqrt{G} \left\{ R - \frac{1}{4} F_{\mu\nu}^I \star G_I^{\mu\nu} - 2g_{A\bar{B}} \partial_\mu z^A \partial^\mu \bar{z}^B - \tilde{g}_{uv} \partial_\mu q^u \partial^\mu q^v \right\} . \quad (1)$$

The n_v complex scalars z^A coming from vector multiplets parameterize a special Kähler manifold K_{n_v} with metric $g_{A\bar{B}}(z, \bar{z})$, whereas the $4n_h$ real scalars q^u a quaternionic manifold Q_{n_h} with metric $\tilde{g}_{uv}(q)$. The gauge field strengths $(F_{\mu\nu}^I, G_{I\mu\nu})$ form a symplectic vector ($I = 0 \dots n_v$) and therefore are not independent

$$G_{I\mu\nu} = \text{Re} \mathcal{N}_{IJ} F_{\mu\nu}^J - \text{Im} \mathcal{N}_{IJ} \star F_{\mu\nu}^J . \quad (2)$$

¹Based on a talk given at the “31st International Symposium Ahrenschoop on the Theory of Elementary Particles” Buckow, September 2-6, 1997.

Like in 10 dimensions we can classify the 4-d supergravity theory in terms brane-solutions (related to elementary and solitonic states); for the 6-d case an analog discussion can be found in [6]. These are the membranes, strings, black holes (0-branes) and instantons (-1-branes). The membranes introduce a mass term and are the analog of the 8-brane solution of massive IIA supergravity. We will ignore this solution furthermore and focus mainly on the other three branes. The string solution is naturally supported by antisymmetric tensors, that couples to their worldvolume. In the action (1) the antisymmetric tensors have been dualized to scalars which are part of the hyper scalars. Black holes couple naturally to vector fields and therefore are solutions with non-trivial vector multiplets. Finally the instantons are dual to the strings and are related to non-trivial hyper scalars. In addition to these counterparts of the 10-d D -branes, we have also purely gravitational solutions, the G -branes as discussed in [7], the wave and Taub-NUT soliton. The branes can also be seen as a classification of the solutions in the number of translational isometries.

These are the “basic” branes of 4-d supergravity. In addition, (threshold) bound states should exist, but so far no solution could be constructed which includes non-trivial vector as well as hyper multiplet scalars. But we can adopt the 10-d duality rules to transform all “basic” solutions into one another.

A good starting point is the black hole or 0-brane. In this case we can neglect the quaternionic part by taking $q^u = \text{constant}$. Let us summarise here only the main points, for details we refer to [3]. The gauge field equations and Bianchi identities are given by $dF^I = dG_I = 0$ and have the solution

$$F_{mn}^I = \frac{1}{2}\epsilon_{mnp}\partial_p\tilde{H}^I \quad , \quad G_{Imn} = \frac{1}{2}\epsilon_{mnp}\partial_p H_I \quad (3)$$

where $m, n, p = 1, 2, 3$ and $(\tilde{H}^I, H_I)$ are harmonic functions. Note, that the timelike components F_{0m}^I and G_{I0m} are fixed in terms of the spatial ones using (2). As next step we need the scalars and the metric, which can be obtained by solving the Killing spinor equations [3]

$$ds^2 = -e^{2U}(dt + \omega_m dx^m)^2 + e^{-2U}dx^m dx^m \quad , \quad z^A = \frac{X^A}{X^0} \quad (4)$$

with

$$\begin{aligned} e^{-2U} &= e^{-K} \equiv i(\bar{X}^I F_I - X^I \bar{F}_I) \quad , \\ \frac{1}{2}e^{2U}\epsilon_{mnp}\partial_n\omega_p &= Q_m \equiv \frac{1}{2}e^{2U}(H_I\partial_m\tilde{H}^I - \tilde{H}^I\partial_m H_I) \end{aligned} \quad (5)$$

and the symplectic section is constraint by

$$-i(X^I - \bar{X}^I) = \tilde{H}^I(x^\mu) \quad , \quad -i(F_I - \bar{F}_I) = H_I(x^\mu) \quad (6)$$

where the harmonic functions $(\tilde{H}^I, H_I)$ are introduced in (3) and $F_I = \frac{\partial}{\partial X^I}F(X)$. This solution is expressed completely in terms of duality invariant quantities, the Kähler potential K and the Kähler connection Q_m . But it is *not* written in a Kähler invariant way, instead we have fixed the Kähler invariance in a proper way. If $\omega_m = 0$ we get the expected static black hole and for special choices for the harmonic

functions we obtain a rotating black hole or generalisations of Taub-NUT and Eguchi-Hanson spaces.

By T -dualizing different directions of the 4-d space time, one obtains the other brane solutions. In order to be concrete let us consider one simple example, which is axion-free, i.e. $\text{Re}z^A = 0$. This can be done by setting $H^0 = H_A = 0$, i.e. the graviphoton is electric and all vector multiplets are magnetic. As prepotential we consider

$$F(X) = \frac{d_{ABC}X^AX^BX^C}{X^0} \quad (7)$$

and the solution is given by [4]

$$e^{-2U} = \sqrt{H_0 d_{ABC}H^AH^BH^C} \quad , \quad z^A = i H_0 H^A e^{2U} \quad (8)$$

with $H_0 = h_0 + q_0/r$, $H^A = h^A + p^A/r$. This black hole is a compactification of the $5 \times 5 \times 5 + \text{mom.}$ intersection, where each of the 5-brane wraps a 4-cycle of the CY_3 and H_0 corresponds to the momentum modes travelling along the common string. A detailed microscopic analysis has been given recently in [2].

In order to obtain the instanton solution we have first to T -dualize the Euclidean time, which is also known as c -map has been used in [5] to map the general solution (5). Note, because the black hole solution considered here, is a type IIA compactification, after this mapping we obtain a type IIB compactification, where the wrapped 5-branes become (instantonic) wrapped 3-branes and the momentum modes are converted into a IIB (-1) -brane, i.e. the intersection is $(-1) \times 3 \times 3 \times 3$. The resulting solution reads [5]

$$\begin{aligned} ds_{inst.}^2 &= e^{-2U} dx^\mu dx^\mu \quad , \quad e^{2\phi} = e^{-2U} \quad , \quad \zeta^0 = \frac{1}{\sqrt{2}} A^0 = \frac{1}{\sqrt{2}H_0} \\ \partial_m \zeta_A &= -\frac{1}{2\sqrt{2}} e^{2U} \text{Im} \mathcal{N}_{AB} \partial_m H^A \quad , \quad z^A \rightarrow q^A \end{aligned} \quad (9)$$

where the first line contains the string-metric and the universal hypermultiplet, i.e. the dilaton ϕ and the former graviphoton gauge potential. In the second line are the scalars coming from the dualization of the magnetic vector fields and also the original vector multiplet scalars. All these scalars group together in $n_v + 1$ hypermultiplets each containing 4 scalars. After having done this dualization one can allow the fields to depend on all 4 coordinates in a symmetric way, i.e. the new harmonic functions are $H_0 = h_0 + q_0/r^2$, $H^A = h^A + p^A/r^2$. Note, the Einstein metric is flat.

As next step we consider the string solution, obtained by T -dualizing one of the spatial direction. This is again a type IIB solution corresponding intersection is $1 \times 5 \times 5 \times 5$

$$\begin{aligned} ds_{string}^2 &= e^{2U} (-dt^2 + dx^2) + e^{-2U} dw d\bar{w} \\ e^{2\phi} &= e^{2U} \quad , \quad \partial_m a \sim \epsilon_{mnp} \partial_p H_0 \quad , \quad \mathcal{H}_A \sim e^{2U} \text{Im} \mathcal{N}_{AB} \star dH^B \quad , \quad \tilde{\mathcal{H}}_A \sim -i e^{2U} g_{AB} \star dz^B \end{aligned} \quad (10)$$

where we choose a parameterization in terms of one complex scalar containing the axion a and the dilaton ϕ . In addition, the compactified D -5-branes and the Hodge

dualization of the scalars give $2n_v$ antisymmetric tensors $(\mathcal{H}_A, \tilde{\mathcal{H}}_A)$. Equivalently, one could dualize them into (hyper) scalars. In doing this T -duality we have to assume a further isometry, i.e. the harmonic functions depend on one coordinate less. Thus the new harmonic functions are general holomorphic functions $H = f(w) + f(\bar{w})$, i.e. they are undetermined from the equations of motion, but restricted due to the proper behaviour at infinity and finite energy, see [8]. Notice however, because of the Kähler structure it is not obvious that the dilaton factorizes in harmonic factors. We will leave a detailed analysis for the future.

One has to keep in mind, that for all these solutions the internal space is fixed (branes wrapped around 4-cycles). Our T -duality transformation acts in the 4-d space time. Duality with respect to the internal manifold would of course not change the 4-d solution, it is given by the duality invariant Kähler potential.

Having these “basic” brane solutions we have to ask for singularities. Let us focus on the instantons and black holes. The harmonic function that we consider here have a pole of $1/r$. As consequence of the constraint for the symplectic section (6) also (X^I, F_I) have a $1/r$ pole. Thus the Kähler potential behaves like $1/r^2$ and the metric is non-singular on the horizon. This property is independent of the number of charges, as long as we keep a non-trivial graviphoton. So, *any generic CY black hole has a non-singular horizon*. This is an important difference to $N = 4, 8$ black holes that need at least 4 charges to get a non-singular horizon. Here, in the CY case, the selfintersection regularize the horizon, which has a throat geometry. Of complete different nature is the instanton solution. Again using the scaling behaviour of the Kähler potential one can show, that for the $r \rightarrow 0$ one reaches a second asymptotic flat region, where the charges and moduli are interchanged. Both regions are connected by wormhole at the minimum of re^{-U} and one of them is strongly coupled ($\phi \rightarrow \infty$), which is very similar analog to the situation in 10 dimensions [9].

Let us end this article with the conformal field theory (CFT) description of the near horizon region of the black hole. Basically it is not necessary to approach the horizon, instead one can scale the physical parameter appropriately (conformal limit). The result is the same – one neglects the constant parts in the harmonic functions in this limit. All non-singular brane configurations are expected to have a super conformal field theory description [10]. As example we consider the solution (8). This is a special case, but if we convince ourselves that it is described by an exact CFT, due to duality also the other charge configurations should be exact. This solution is a compactified 5-d magnetic string [4]. Near the horizon, the spherical part becomes constant and a reduction over this spherical part yields a (negative) cosmological constant giving an AdS_3 geometry. By proper boundary conditions a conformal algebra is realized as diffeomorphism group at spatial infinity [11] (this is also nicely summarized in [12]) and the corresponding solution is the BTZ black hole [13] (for a discussion of the entropy and duality see especially [14, 12, 15]).

In the following we will sketch this procedure for our solution, a more detailed

analysis will be given elsewhere. Near the horizon the magnetic string becomes [4]

$$ds^2 = \frac{r}{D^{1/3}} \left[-dt^2 + dx^2 + \frac{q_0}{r} (dx - dt)^2 \right] + D^{2/3} \left(\frac{dr}{r} \right)^2 + D^{2/3} d\Omega_2, \quad (11)$$

where $D = d_{ABC} p^A p^B p^C$. As next step one compactifies the S^2 part. The 5-d sugra action contains besides the Einstein-Hilbert term, a gauge field part and a scalar part. Near the horizon the scalar part drops out and the gauge field part contributes to the cosmological constant $G_{IJ} F^I \cdot F^J = D^{-2/3}$ with $G_{IJ}(X)$ as the 5-d gauge coupling matrix. After this reduction the 3-d Einstein metric reads

$$ds_E^2 = D r \left[-dt^2 + dx^2 + \frac{q_0}{r} (dx - dt)^2 \right] + D^2 \left(\frac{dr}{r} \right)^2 \quad (12)$$

or in other coordinates ($\rho^2 = D(r + q_0)$, $\tau = 2Dt$ and $x = \phi$)

$$ds_E^2 = -\frac{(\rho^2 - q_0 D)^2}{4\rho^2 D^2} d\tau^2 + \rho^2 \left(d\phi - \frac{q_0}{2\rho^2} d\tau \right)^2 + \frac{4D^2 \rho^2}{(\rho^2 - q_0 D)^2} d\rho^2. \quad (13)$$

This is the extreme 3-d BTZ black hole [13], with the angular momentum $J = q_0$, the mass $M = \frac{q_0}{2D}$ and the cosmological constant $\Lambda^{-1} \sim l = 2D$. Thus excited momentum modes along the 5-d string correspond to angular momentum for this BTZ black hole. As it has been shown in [16], 3-d AdS gravity can be written as a Chern-Simons theory and furthermore, due to the cylindric symmetry ($\phi \simeq \phi + 2\pi$) it reduces to a sum of two $SL(2, \mathbb{R})$ WZW theories for the left and right moving modes (in the BPS limit only one sector survive). Finally, implementing the boundary condition at spatial infinity we end up with a Liouville theory living on the boundary [17]. So, summarizing all steps we have the sequence for the actions

$$S_{sugra_5} \rightarrow S_{AdS_3} \sim k S_{CS} \sim k S_{WZW} \sim k S_L \quad (14)$$

where up to numerical factors, the central charge of the CFT is given by $k \sim \frac{D}{G_N}$, where we introduced the 3-d Newton constant G_N to get a dimensionless quantity.

Let us conclude with implications of this CFT for the Matrix description. It has been suggested recently that the matrix limit of M -theory compactified on CY_3 is given by the conifold point, which is mirror dual to the vanishing 6-cycle [18]. An important point in the matrix limit is to ensure the decoupling of massless states (decoupling from the bulk). For a T^6 compactification this is in general not the case (although as discussed in [19] there is may be a loophole by a gauged world volume theory). In order to discuss this question here we consider only 5-branes which wrap vanishing cycles. Notice, in our setup we approaching the conifold point from the other side, i.e. we consider vanishing 4-cycles. As consequence, the corresponding 4-d black hole becomes massless [20] or, for the magnetic sting, $D \rightarrow 0$ in this limit (note, if L counts the vanishing cycles $X^L \sim p^L \rightarrow 0$ at this point). As consequence the central charge of the CFT vanish, which coincides with the expectation that at the transition point the worldvolume theory is scale invariant. But more important,

in this limit the boundary degrees of freedom (encoded in the Liouville action S_L) decouple, like the Liouville mode in critical string theory. This supports the idea that the matrix description for M -theory compactified on CY_3 [18] yields an consistent picture.

If one goes away from the conformal point one has to take into account the additional boundary modes and an immediate question is to ask for their contribution to the black hole entropy of (8). Of course, quantizing the boundary CFT one obtains a shift in the central charge (from the Liouville mode). This is equivalent to a shift in the cosmological constant $D \rightarrow D + const.$, or the corrected black hole entropy reads

$$\mathcal{S} \sim 2\pi \sqrt{q_0(d_{ABCP}{}^A p^B p^C + const.)} . \quad (15)$$

It would be very interesting to discuss these corrections in more detail. Note this shift is independent of the magnetic charges that parameterize the the 4-cycles of the CY. Similar shifts of D have recently been discussed in terms of higher loop corrections for the black hole entropy [2, 21], but the corrections there are linear in the magnetic charges. Because the shift here is not sensitive to any internal CY structure, one should expect that it is a pure supergravity effect, i.e. related to terms in the prepotential that contain only X^0 . A further interesting line of continuation would be to keep the scalar fields as matter part in this CFT description.

Acknowledgements

I am grateful I. Brunner, G. Cardoso, S. Kachru, A. Karch and R. Schimmrigk for helpful discussions. This work is support by the DFG.

References

- [1] S. Ferrara, R. Kallosh and A. Strominger, *N=2 extremal black holes*, hep-th/9508072; K. Behrndt, G. Lopes-Cardoso, B. de Wit, R. Kallosh, D. Lüst and T. Mohaupt, *Classical and supersymmetric N=2 black holes*, hep-th/9610105.
- [2] J. Maldacena, A. Strominger and E. Witten, *Black hole entropy in M-theory*, hep-th/9711053; C. Vafa, *Black holes and Calabi-Yau threefolds* hep-th/9711067.
- [3] K. Behrndt, D. Lüst and W.A. Sabra, *Stationary solutions of N=2 supergravity*, hep-th/9705169.
- [4] K. Behrndt, *Quantum corrections for D = 4 black holes and D = 5 strings*, hep-th/9610232.
- [5] K. Behrndt, I. Gaida, D. Lüst, S. Mahapatra and T. Mohaupt, *From type IIA black holes to T dual type IIB D instantons in N=2, D = 4 supergravity*, hep-th/9706096.

- [6] K. Behrndt, E. Bergshoeff and B. Janssen, *Intersecting d-branes in ten-dimensions and six-dimensions*, [hep-th/9604168](#) .
- [7] C. Hull *Gravitational duality, branes and charges*, [hep-th/9705162](#).
- [8] B.R. Greene, A. Shapere, C. Vafa and S.-T. Yau, *Nucl. Phys.* **B337** (1990) 1, M. Bourdeau and G. Lopes Cardoso, *Finite energy solutions in three-dimensional heterotic string theory*, [hep-th/9709174](#).
- [9] G.W. Gibbons, M.B. Green and M.J. Perry, *Instantons and 7-branes in type IIB superstring theory*, [hep-th/9511080](#).
- [10] P. Claus, R. Kallosh and A. Van Proeyen, *M five-brane and superconformal (0,2) tensor multiplet in six-dimensions*, [hep-th/9711161](#); J. Maldacena, *The Large N limit of superconformal field theories and supergravity*, [hep-th/9711200](#); R. Kallosh, J. Kumar and A. Rajaraman, *Special conformal symmetry of worldvolume action*, [hep-th/9712073](#); S. Ferrara and C. Fronsda, *Conformal Maxwell theory as a singleton field theory on $adS(5)$, IIB three-branes and duality*, [hep-th/9712239](#).
- [11] J.D. Brown and M. Henneaux, *Comm. Math. Phys.* **104** (1986) 207.
- [12] A. Strominger, *Black hole entropy from near horizon microstates*, [hep-th/9712251](#).
- [13] M. Banados, C. Teitelboim and J. Zanelli, *The Black hole in three-dimensional space-time*, [hep-th/9204099](#); M. Banados, M. Henneaux, C. Teitelboim and J. Zanelli, *Geometry of the (2+1) black hole*, [gr-qc/9302012](#).
- [14] S. Carlip, *The Statistical mechanics of the (2+1)-dimensional black hole*, [gr-qc/9409052](#).
- [15] S. Hyun, *U duality between three-dimensional and higher dimensional black holes*, [hep-th/9704005](#); K. Sfetsos and K. Skenderis, *Microscopic derivation of the Bekenstein-Hawking entropy formula for nonextremal black holes*, [hep-th/9711138](#); H.J. Boonstra, B. Peeters and K. Skenderis, *Duality and asymptotic geometries*, [hep-th/9706192](#).
- [16] A. Achucarro and P.K. Townsend, *Phys. Lett.* **B180** (1986) 89; E. Witten, *Nucl. Phys.* **B311** (1988) 46.
- [17] O. Coussaert, M. Henneaux and P. van Driel, *The Asymptotic dynamics of 3-d Einstein gravity with a negative cosmological constant*, [gr-qc/9506019](#)
- [18] S. Kachru, A. Lawrence and E. Silverstein, *On the matrix description of Calabi-Yau compactifications*, [hep-th/9712223](#).
- [19] K. Behrndt, *On central charges and the entropy in matrix theory*, [hep-th/9710228](#).

- [20] A. Strominger, *Massless black holes and conifolds in string theory*, hep-th/9504090; K. Behrndt, D. Lüster and W.A. Sabra, *Moving moduli, Calabi-Yau phase transitions and massless BPS configurations in type II superstrings*, hep-th/9708065
- [21] K. Behrndt, G. Lopes Cardoso, B. de Wit, D. Lüster, T. Mohaupt and W. Sabra, *Higher-order black hole solutions in $N=2$ supergravity and Calabi-Yau string backgrounds*, hep-th/9801081.